

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human Niemann-Pick Type C1/NPC1 in direct ELISAs. Detects human, mouse, and rat Niemann-Pick Type C1/NPC1 in Western blots.
Source	Monoclonal Rabbit IgG ₁ Clone # 1318A
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Synthetic peptide containing Human Niemann-Pick Type C1/NPC1
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Knockout Validated	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NPC intracellular cholesterol transporter 1 or Niemann-Pick C1 protein (NPC1) is a 1,278 aminoacids (aa) intracellular cholesterol transporter which plays an important role in cholesterol transport from the endosomal/lysosomal compartment. NPC1 was identified as the gene that when mutated, results in Niemann-Pick disease, type C, a rare neurovisceral lipid storage disorder resulting from autosomal recessively inherited loss-of-function mutations in either NPC1 or NPC2. This disrupts intracellular lipid transport, leading to the accumulation of lipid products in the late endosomes and lysosomes. Approximately 95% of NPC patients are found to have mutations in the NPC1 gene. In humans, at least one other isoform, missing aa 519-586 is known. Human NPC1 protein sequence is 87% identical to both, mouse and rat NPC1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.