

**DESCRIPTION**

|                           |   |
|---------------------------|---|
| <b>Species Reactivity</b> | Human   |
| <b>Specificity</b>        | Detects human CD44 in direct ELISA.   |
| <b>Source</b>             | Monoclonal Mouse IgG <sub>2B</sub> Clone # 1058722  |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant                                    |
| <b>Immunogen</b>          | Chinese Hamster Ovary cell line, CHO-derived human CD44<br>Gln21-Thr222<br>Accession # P16070 |
| <b>Conjugate</b>          | Alexa Fluor 488<br>Excitation Wavelength: 488 nm<br>Emission Wavelength: 515-545 nm           |
| <b>Formulation</b>        | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide                                  |

\*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

**APPLICATIONS**

|   |  |
|---|--|
| <b>Please Note:</b> Optimal dilutions should be determined by each laboratory for each application. <i>General Protocols</i> are available in the Technical Information section on our website. |  |
| <b>Immunocytochemistry</b>  | Optimal dilution of this antibody should be experimentally determined. |
| <b>Immunohistochemistry</b>   | Optimal dilution of this antibody should be experimentally determined. |

**PREPARATION AND STORAGE**

|                                |   |
|--------------------------------|---|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

**BACKGROUND**

CD44 is a ubiquitously expressed protein that is the major receptor for hyaluronan and exerts control over cell growth and migration (1-3). Human CD44 has a 20 amino acid (aa) signal sequence, an extracellular domain (ECD) with a 100 aa hyaluronan-binding disulfide-stabilized link region and a 325-530 aa stem region, a 21 aa transmembrane domain, and a 72 aa cytoplasmic domain. CD44 transcripts undergo complex alternative splicing, and, within the stem, ten variably spliced exons (v1-10 corresponding to exons 6-15; although human CD44 lacks v1/exon 6) produce multiple protein isoforms (1-4). The standard or hematopoietic form, CD44H, does not include the variable segments (1-4). Cancer aggressiveness and T cell activation have been correlated with expression of specific isoforms (1, 4, 5). Human CD44v5 contains exon 10 and is associated with tumor progression and metastasis in many types of cancer including breast, colon, lung, renal, skin, and ovarian tumors (6, 12). With variable N- and O-glycosylation and splicing within the stalk, CD44 can range from 80 to 200 kDa (1). Within the N-terminal invariant portion of the ECD (aa 21-222), human CD44 shares 76%, 76%, 86%, 83% and 79% identity with corresponding mouse, rat, equine, canine and bovine CD44, respectively. The many reported functions of CD44 fall within three categories (1). First, CD44 binds hyaluronan and other ligands within the extracellular matrix and can function as a "platform" for growth factors and metalloproteinases. Second, CD44 can function as a co-receptor that modifies activity of receptors including MET and the ERBB family of tyrosine kinases. Third, the CD44 intracellular domain links the plasma membrane to the actin cytoskeleton via the ERM proteins, ezrin, radixin and moesin. CD44 can be synthesized in a soluble form (6) or may be cleaved at multiple sites by either membrane-type matrix metalloproteinases, or ADAM proteases to produce soluble ectodomains (8, 9). The cellular portion may then undergo gamma secretase-dependent intramembrane cleavage to form an A beta-like transmembrane portion and a cytoplasmic signaling portion that affects gene expression (10, 11). These cleavage events are thought to promote metastasis by enhancing tumor cell motility and growth (1, 8). CD44v5 plays important role in cancer progression involving in cell migration and invasion and is considered a functional cancer biomarker (6, 12).

**PRODUCT SPECIFIC NOTICES**

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.