

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IFN-alpha 4 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse IFN-1, -2, -6, -7, -9, -11, -12, -13, -15 or -16 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 1020125
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell HEK293-derived mouse IFN-alpha 4 Leu23-Glu186 Accession # P07351
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The interferons (IFN) are a family of cytokines with potent antiviral, antiproliferative and immunomodulatory properties, and are classified based on their binding specificity to cell surface receptors (1). The type I IFNs bind to the interferon alpha receptor (IFNAR), which consists of two subunits: IFNAR1 (α-subunit) and IFNAR2 (β-subunit). This binding contributes to TNF-alpha induced signaling (2, 3). Both the human and mouse genome code for more than a dozen closely related IFN-alpha subtypes and the various IFN-alphas share about 80% sequence homology among them (4, 5). The mouse IFN-alpha 4 consists of 186 amino acids (aa) including a 24 aa signal peptide and a 162 aa IFN-alpha 4 mature domain. The mature mouse IFN-alpha 4 shares 58% and 76% sequence identity with mature human and rat IFN-alpha 4, respectively. The mouse IFN-alpha 4 reduces the local replication of murine cytomegalovirus in the tibia muscle (6). mIFN-A4 is a strong activator of Mx gene and was shown to be very effective in reducing splenomegaly (7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.