

## Human EBI2 Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG<sub>2A</sub> Clone # 486633 Catalog Number: FAB10272G

100 µg

| DESCRIPTION        |   |  |
|--------------------|---|--|
| Species Reactivity | Human   |  |
| Specificity        | Detects human EBI2 in direct ELISAs.  |  |
| Source             | Monoclonal Mouse IgG <sub>2A</sub> Clone # 486633   |  |
| Purification       | Protein A or G purified from hybridoma culture supernatant  |  |
| Immunogen          | Mouse myeloma cell line NS0-derived transfected withrecombinant human EBI2 Accession # P32249   |  |
| Conjugate          | Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm   |  |
| Formulation        | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.  |  |
|                    | *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |  |

| APPLICATIONS  |                                 |   |  |
|---|---------------------------------|---|--|
| Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website. |                                 |   |  |
|   | Recommended<br>Concentration    | Sample  |  |
| Flow Cytometry  | 0.25-1 μg/10 <sup>6</sup> cells | HEK293 Human Cell Line Transfected with Human EBI2 and eGFP and A549 human lung carcinoma cell line |  |

| PREPARATION AND STORAGE                                |   |  |
|--|---|--|
| Shipping   | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |  |
| Stability & Storage Protect from light. Do not freeze. |   |  |
|  | <ul> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>  |  |

## BACKGROUND

Epstein-Barr virus-induced G-protein coupled receptor 2, also known as EBI2 or GPR183 is a G-protein coupled receptor (GPCR) expressed in lymphocytes that acts as a chemotactic receptor for B-cells, T-cells, splenic dendritic cells, and monocytes/macrophages. The GPR183 gene was identified by the up-regulation of its expression upon Epstein-Barr virus infection of the Burkitt's lymphoma cell line BL41. GPR183 collaborates with CXCR5 to mediate B cell homing within a lymph node, probably by forming a heterodimer with CXCR5 that affects its interaction with CXCL13

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

