

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CTRP1/C1qTNF1 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 395520
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived human CTRP1/C1qTNF1 Arg26-Pro281 Accession # Q9BXJ1
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

C1qTNF1 (CTRP1) is an approximately 35 kDa member of the C1q family of secreted proteins and plays a role in energy metabolism and inflammation (1, 2). C1qTNF1 contains a collagen-like region and one C1q-like domain (3). Mature human C1qTNF1 shares 80% aa sequence identity with mouse and rat C1qTNF1. Circulating levels of C1qTNF1 are elevated in obesity, hypertension, and diabetes but can be decreased in the serum of diet-induced obese mice (4-6). C1qTNF1 expression is up-regulated in atherosclerotic plaques or adipose tissue by oxidized LDL or inflammatory cytokines (3, 7, 8). In turn, it induces the expression of inflammatory cytokines (7, 9) and the up-regulation of adhesion proteins on vascular endothelial cells (8). Systemically administered C1qTNF1, in contrast, can limit tissue damage following myocardial infarction (9). In skeletal muscle, C1qTNF1 promotes fatty acid oxidation, energy expenditure, insulin sensitivity, and glucose uptake and glycolysis (6, 10). It also induces the proliferation of immature chondrocytes (11) and aldosterone synthesis in the adrenal cortex (4). R&D Systems in-house testing indicates that C1qTNF1 binds to BAI3, consistent with the reported interactions between BAI3 and C1qL proteins (12).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.