

DESCRIPTION

Species Reactivity	Human
Specificity	Detects recombinant human EN-RAGE in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 161205
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human EN-RAGE Met1-Glu92 Accession # P80511
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human whole blood

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

EN-RAGE, also known as S100A12 and Calgranulin C, is a 10 kDa member of the S100 (soluble in 100% saturated ammonium sulfate) family of EF-hand calcium-binding proteins. Like other S100 proteins, S100A12 is small and generally acidic (1-3). EN-RAGE forms noncovalent homodimers in the absence of divalent cations (4-6); calcium and zinc promote the formation of higher order assemblies including tetramers and hexamers (4, 7, 8). The ability of S100A12 to chelate zinc enables it to inhibit the zinc-dependent metalloproteases MMP-2, -3, and -9 (8). S100A12 also forms heterodimers with S100A9 and binds to RAGE (Receptor for Advanced Glycation End-products), Annexin V, and several cytosolic enzymes involved in energy metabolism (9, 10). The hexameric form of EN-RAGE in particular binds RAGE with high affinity (7). EN-RAGE induces a variety of inflammatory responses including the *in vivo* recruitment of neutrophils, monocytes, and mast cells and the activation of mast cells and vascular endothelial cells (9, 11-14). EN-RAGE is found at elevated levels under inflammatory conditions such as asthma, gout, rheumatoid arthritis synovial fluid, and atherosclerosis (8, 12, 14). S100A12 also promotes neurite outgrowth in isolated hippocampal neurons (15). An ortholog of S100A12 has not been identified in rodents, but the human protein is functional in mice and rats (11-16).

References:

1. Santamaria-Kisiel, L. *et al.* (2006) *Biochem. J.* **396**:201.
2. Leclerc, E. *et al.* (2009) *Biochim. Biophys. Acta* **1793**:993.
3. Wicki, R. *et al.* (1996) *Cell Calcium* **20**:459.
4. Moroz, O.V. *et al.* (2009) *BMC Biochem.* **10**:11.
5. Miranda, L.P. *et al.* (2001) *FEBS Lett.* **488**:85.
6. Vogl, T. *et al.* (1999) *J. Biol. Chem.* **274**:25291.
7. Xie, J. *et al.* (2007) *J. Biol. Chem.* **282**:4218.
8. Goyette, J. *et al.* (2009) *J. Immunol.* **183**:593.
9. Hatakeyama, T. *et al.* (2004) *Eur. J. Biochem.* **271**:3765.
10. Hofmann, M.A. *et al.* (1999) *Cell* **97**:889.
11. Yang, Z. *et al.* (2001) *J. Leukoc. Biol.* **69**:986.
12. Rouleau, P. *et al.* (2003) *Clin. Immunol.* **107**:46.
13. Yan, W.X. *et al.* (2008) *J. Biol. Chem.* **283**:13035.
14. Yang, Z. *et al.* (2007) *J. Allergy Clin. Immunol.* **119**:106.
15. Mikkelsen, S.E. *et al.* (2001) *J. Neurochem.* **79**:767.
16. Fuellen, G. *et al.* (2004) *OMICS* **8**:334.

Human EN-RAGE/S100A12 Alexa Fluor® 750-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 161205

Catalog Number: FAB10521S
100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.