

DESCRIPTION

Species Reactivity	Cynomolgus Monkey
Specificity	Detects Cynomolgus monkey IL-23R in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 1030030
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell HEK293-derived Cynomolgus monkey IL-23R Gly24-Asp353 Accession # XP_005543141.1
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 23 (IL-23) is a heterodimeric cytokine composed of two disulfide-linked subunits, a p19 subunit that is unique to IL-23, and a p40 subunit that is shared with IL-12 (1-5). The functional IL-23 receptor complex consists of two receptor subunits, the IL-12 receptor beta 1 subunit (IL-12R beta 1) and the IL-23-specific receptor subunit (IL-23R) (3). Human IL-23R cDNA encodes a 629 amino acids (aa) type I transmembrane protein with a 23 aa residue signal peptide, a 330 aa residue extracellular domain, a 23 aa residue transmembrane domain and a 253 aa residue cytoplasmic region. IL-23 R shares structural features with the IL-12R beta 2, including an N-terminal Ig-like domain, two cytokine receptor domains and multiple glycosylation sites in the extracellular domain. IL-23R lacks the three extracellular membrane-proximal fibronectin-type III domains present on IL-12R beta 2. IL-23R has a WQPWS sequence in the transmembrane-proximal cytokine receptor domain similar to the cytokine receptor signature WSXWS motif (6). The cytoplasmic region of IL-23R has three potential Src homology 2 domain-binding sites and two potential Stat-binding sites. The gene for human IL-23R is located on human chromosome 1 within 150 kb of IL-12R beta 2. Based on quantitative real-time PCR, human IL-23R mRNA is expressed in a human Th1 and Th0 clone as well as several NK cell lines and clones. Low but detectable levels of IL-23R mRNA is also expressed in EBV-transformed B cells and activated PBMC. IL-23 initiates a signal transduction cascade similar to that of IL-12, and involves Jak2, Tyk2, Stat1, Stat3, Stat4, and Stat5 (2). The Cynomolgus IL-23R shares 96%, 71% and 77% amino acid sequence identity to Human, mouse, and rat IL-23R, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.