

## DESCRIPTION

|   |   |
|---|---|
| <b>Species Reactivity</b>   | Human   |
| <b>Specificity</b>  | Detects human CD3ε in direct ELISAs.  |
| <b>Source</b>   | Monoclonal Mouse IgG <sub>2B</sub> Clone # 1033713  |
| <b>Purification</b>   | Protein A or G purified from hybridoma culture supernatant                                  |
| <b>Immunogen</b>  | Human embryonic kidney cell HEK293-derived human CD3ε<br>Gln22-Asp126<br>Accession # P07766 |
| <b>Conjugate</b>  | Alexa Fluor 750<br>Excitation Wavelength: 749 nm<br>Emission Wavelength: 775 nm             |
| <b>Formulation</b>  | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide                                |
| *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |   |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

|                             |  |
|-----------------------------|--|
| <b>Western Blot</b>         | Optimal dilution of this antibody should be experimentally determined. |
| <b>Immunohistochemistry</b> | Optimal dilution of this antibody should be experimentally determined. |

## PREPARATION AND STORAGE

|                                |   |
|--------------------------------|---|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

## BACKGROUND

CD3ε (T-Cell Surface Glycoprotein CD3 Epsilon) is a type I transmembrane protein that belongs to the Ig (Immunoglobulin) superfamily. Human CD3ε consists of 104 amino acid (aa) extracellular domain, 26 aa transmembrane domain and 55 aa cytoplasmic domain (1). With the extracellular domain, human CD3ε shares 44.2% and 49% homology with mouse and rat CD3ε respectively. There are four CD3 proteins: CD3δ, CD3ε, CD3γ, and CD3ζ. All CD3 proteins contain ITAM (Immunoreceptor Tyrosine-based Activation Motifs) in the cytoplasmic tail, which becomes phosphorylated by Src family protein tyrosine kinases LCK and FYN upon TCR engagement (2). CD3 proteins form heterodimers of CD3δ/CD3ε and CD3γ/CD3ε, which bind to TCR and form trimeric TCRα/CD3ε/CD3γ and TCRβ/CD3γ/CD3ε. The resulting heterohexamer further associates with CD3ζ homodimer and forms TCR/CD3 signaling complex (3, 4). Similar complex is also formed with TCRγ and TCRδ (5). CD3ε plays essential role in adaptive immune response. Mutations in CD3ε lead to SCID (severe combined immunodeficiency) (6).

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.