

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PGC1α in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 474915
Purification	Protein A or G purified
Immunogen	<i>E. coli</i> -derived human PGC1α Glu11-Ile280 Accession # Q9UBK2
Conjugate	Alexa Fluor Plus 555 Excitation Wavelength: 558 nm Emission Wavelength: 572 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

DATA

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PGC1α (PPAR-γ Coactivator 1 alpha; also LEM6) is a 97-120 kDa member of the PGC-1 family of proteins. It is expressed in select cell types, including brown adipocytes, skeletal muscle and hepatocytes. PGC1α participates in both RNA processing and transcriptional coactivation in conjunction with multiple nuclear hormone receptors such as PPARγ, RAR and TR. Human PGC1α is 798 amino acids (aa) in length. It contains an LxxLL nuclear receptor binding motif (aa 144-148), one PPAR-γ interaction domain (aa 293-339), two NLSs and an RNA binding/processing region (aa 566-710). PGC1α activity is regulated by phosphorylation. AMPK is known to phosphorylate Thr178 and Ser539, promoting cotranscriptional activity. Conversely, Akt-mediated phosphorylation at Ser571 is reported to downregulate PGC1α activity. This latter effect is achieved by an initial Ser571 phosphorylation, followed by GCN5 binding and subsequent PGC1α acetylation that promotes PGC-1α dissociation from target gene promoters.

PRODUCT SPECIFIC NOTICES

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.