

Human PD-1 Alexa Fluor® 700-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 1026519
Catalog Number: FAB10866N
100 µg

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PD-1 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 1026519
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Synthetic peptide containing human PD-1
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.

*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	HEK293 Human Cell Line transfected with Human PD-1 and eGFP

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Programmed Death-1 receptor (PD-1), also known as CD279, is type I transmembrane protein belonging to the CD28 family of immune regulatory receptors (1). Other members of this family include CD28, CTLA-4, ICOS, and BTLA (2-5). Mature human PD-1 consists of a 148 amino acid (aa) extracellular region (ECD) with one immunoglobulin-like V-type domain, a 24 aa transmembrane domain, and a 95 aa cytoplasmic region. The human PD-1 ECD shares 65% aa sequence identity with the mouse PD-1 ECD. The cytoplasmic tail contains two tyrosine residues that form the immunoreceptor tyrosine-based inhibitory motif (ITIM) and immunoreceptor tyrosine-based switch motif (ITSM) that are important for mediating PD-1 signaling. PD-1 acts as a monomeric receptor and interacts in a 1:1 stoichiometric ratio with its ligands PD-L1 (B7-H1) and PD-L2 (B7-DC) (6, 7). PD-1 is expressed on activated T cells, B cells, monocytes, and dendritic cells while PD-L1 expression is constitutive on the same cells and also on nonhematopoietic cells such as lung endothelial cells and hepatocytes (8, 9). Ligand of PD-L1 with PD-1 induces co-inhibitory signals on T cells promoting their apoptosis, anergy, and functional exhaustion (10). Thus, the PD-1: PD-L1 interaction is a key regulator of the threshold of immune response and peripheral immune tolerance (11). Finally, blockade of the PD-1: PD-L1 interaction by either antibodies or genetic manipulation accelerates tumor eradication and shows potential for improving cancer immunotherapy (12, 13, 14).

References:

1. Ishida, Y. *et al.* (1992) EMBO J. **11**:3887.
2. Sharpe, A.H. and G. J. Freeman (2002) Nat. Rev. Immunol. **2**:116.
3. Coyle, A. and J. Gutierrez-Ramos (2001) Nat. Immunol. **2**:203.
4. Nishimura, H. and T. Honjo (2001) Trends Immunol. **22**:265.
5. Watanabe, N *et al.* (2003) Nat. Immunol. **4**:670.
6. Zhang, X. *et al.* (2004) Immunity **20**:337.
7. Lázár-Molnár, E. *et al.* (2008) Proc. Natl. Acad. Sci. USA **105**:10483.
8. Nishimura, H *et al.* (1996) Int. Immunol. **8**:773.
9. Keir, M.E. *et al.* (2008) Annu. Rev. Immunol. **26**:677.
10. Butte, M.J. *et al.* (2007) Immunity **27**:111.
11. Okazaki, T. *et al.* (2013) Nat. Immunol. **14**:1212.
12. Iwai, Y. *et al.* (2002) Proc. Natl. Acad. Sci. USA **99**: 12293.
13. Nogrady, B. (2014) Nature **513**:S10.
14. Swaika, A. *et al.* (2015) Mol. Immunol. **67**: 4

Human PD-1 Alexa Fluor® 700-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 1026519

Catalog Number: FAB10866N

100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.