

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human EDA2R/TNFRSF27/XEDAR in ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) 41BB, rhBAFF R, rhCD27, rhCD30, rhCD40, rhDR-3, rhDR-6, rhEDAR, rhFAS, rhGITR, rhHVEM, rhLTR-b, rhNGFR, rhOPG,
Source	Monoclonal Mouse IgG ₁ Clone # 173103
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human EDA2R/TNFRSF27/XEDAR Met1-Glu136 Accession # Q9HAV5
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

X-linked Ectodysplasin Receptor (XEDAR), also known as EDA2R and TNFRSF27, is an approximately 45 kDa transmembrane protein in the TNF receptor superfamily (1). Mature human XEDAR consists of a 136 amino acid (aa) extracellular domain (ECD), a 21 aa transmembrane segment, and a 140 aa cytoplasmic domain (2). Within the ECD, human XEDAR shares 87% aa sequence identity with mouse and rat XEDAR. A 55 kDa long isoform of human XEDAR carries a 21 aa insertion in the juxtamembrane cytoplasmic domain (3). A 20 kDa fragment of the ECD can be shed by metalloprotease mediated cleavage (4). XEDAR binds selectively to the EDA-A2 variant of Ectodysplasin (EDA), while the closely related receptor EDAR binds selectively to the EDA-A1 variant (2). Other than a 2 aa deletion in its TNF-like domain, EDA-A2 is identical to EDA-A1 (2). Mutations in both EDAR and EDA are associated with hypohidrotic ectodermal dysplasia (HED), a disorder of hair, tooth, and eccrine sweat gland morphogenesis (5). XEDAR itself is strongly associated with androgenetic alopecia (male hair loss) (6). XEDAR is widely expressed, notably in embryonic basal epidermal cells and maturing hair follicles (2, 7, 8). Even though it does not contain a cytoplasmic death domain, XEDAR can associate with Fas and induce EDA-A2 dependent apoptosis (7, 9). Its transcription is directly induced by p53, and XEDAR mediated cell death is p53 dependent (7, 10). XEDAR is down-regulated in breast, colon, and lung cancers, particularly in cases with p53 mutations (7, 11). XEDAR also plays a role in EDA-A2 induced skeletal muscle degeneration and osteoblast differentiation (8, 12).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.