

Mouse TROP-2 Fluorescein-conjugated **Antibody**

Antigen Affinity-purified Polyclonal Goat IgG Catalog Number: FAB1122F


HDCI.	
	100 ua

DESCRIPTION		
Species Reactivity	Mouse	
Specificity	Detects mouse TROP-2 in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant humar TROP-2 is observed.	
Source	Polyclonal Goat IgG	
Purification	Antigen Affinity-purified	
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse TROP-2 Ser82-Thr268 Accession # Q8BGV3	
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)	
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 μg/10 ⁶ cells	See Below



Detection of TROP-2 in mIMCD-3 Mouse Epithelial Cell Line by Flow Cytometry. mIMCD-3 mouse epithelial cell line was stained with Goat Anti-Mouse TROP-2 Fluorescein-conjugated Antigen Affinitypurified Polyclonal Antibody (Catalog # FAB1122F, filled histogram) or isotype control antibody (Catalog # IC108F, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

TROP-2, also named tumor-associated calcium signal transducer 2 (TACSTD2), GA733 tumor associated antigen, and epithelial glycoprotein-1 (EGP-1), is a type I transmembrane protein highly expressed in carcinomas. It was originally identified as an antigen present on human gastrointestinal tumors and is the second of two members of this family. The other family member is GA7332, also called EpCAM, TROP1, 171A, gp40 and KSA. TROP2 can transduce an intracellular calcium signal and may play a role in tumor growth. It also has adhesive functions.

Rev. 2/6/2019 Page 1 of 1

