

**DESCRIPTION**


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse TER-119 in Western blots. This antibody has been shown to react with cells of the erythroid lineage in embryonic yolk sac, fetal liver, adult bone marrow, and adult peripheral blood.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # TER-119
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	C57BL/6 mouse day-14 fetal liver cells
<b>Conjugate</b>	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	5 µL/10 <sup>6</sup> cells	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> ● 12 months from date of receipt, 2 to 8 °C as supplied.

**BACKGROUND**

The monoclonal antibody TER-119 is isolated from a hybridoma generated using splenocytes from a rat subcutaneously injected with day 14 BALB/c fetal liver cells (1). The TER-119 monoclonal antibody reacts with erythroid cells from the early proerythroblast to mature erythrocyte stages of development (1). The 52 kDa ligand for TER-119 is associated with glycoprotein A on erythrocytes (1). TER-119 antibodies are frequently used in combination with other lineage depletion antibodies to enrich for mouse hematopoietic stem cells (2, 3).

**References:**

1. Kina, T. *et al.* (2000) Br. J. Haematol. **109**:280.
2. Ikuta, K. *et al.* (1990) Cell **62**:863.
3. Osawa, M.Y. *et al.*, (1996) Hematopoietic Stem Cells in *Weir's Handbook of Experimental Immunology*, Vol. 2, 5th Edition. Herzenberg, L.A. *et al.* eds. Blackwell Science, Cambridge, MA. pp. 66.1-66.5

**PRODUCT SPECIFIC NOTICES**

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.