

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human FGFR1 alpha (IIIc) in direct ELISA.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 1058809
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Human embryonic kidney cell, HEK293-derived human FGFR1 alpha Arg22-Glu376 Accession # P11362.3
<b>Conjugate</b>	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

<b>Flow Cytometry</b>	Titration recommended for optimal concentration with starting range of 0.1-1 µg/1 million cells. Sample used for this experiment was HEK293 cells transfected with hFGFR1 and eGFP.
-----------------------	---

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Fibroblast growth factor receptor 1 (FGFR1) belongs to a family of type I transmembrane tyrosine kinases which mediate the biological functions of FGFs that are involved in a multitude of physiological and pathological cellular processes (1). The FGFR family is comprised of 4 structurally conserved members (FGFR1-4) all possessing an extracellular domain (ECD) with three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and cytoplasmic split tyrosine-kinase domain (1, 2). The ECD of mature, full-length FGFR1 shares 98% amino acid sequence identity with mouse FGFR1. Alternative splicing generates multiple forms of FGFR1-3, each with unique signaling characteristics (1-3). For FGFR1, alternative splicing of the ECD generates FGFR1A, FGFR1B, and FGFR1G isoforms of the receptor with the A isoform containing three Ig domains, while the B and G isoforms lack the N-terminal IgI domain (3). Additional splicing of the IgIII domain, results in IIIa, IIIb, or IIIc isoforms (3). Only the alpha isoform has been identified for FGFR3 and FGFR4 and FGFR4 also lacks the IIIb and IIIc splicing events (4). The FGFR splice variants also exhibit distinct and varying binding affinities for different FGF ligands (2). FGFRs mediate the FGF signaling cascade which regulate developmental processes including cellular proliferation, differentiation, and migration, morphogenesis, and patterning (5). FGFRs transduce the signals through three dominant pathways including RAS/MAPK, PI3K/AKT, and PLCγ (6). FGFR1 the most abundant FGFR and is widely expressed in many adult human tissues, but the splice variants display distinct tissue-specific differences with IIIc splice variants expressed in mesenchymal tissue (4, 7, 8). Mutations in FGFR1 or misregulation of FGFR1 mediated signaling is found in multiple diseases, with FGFR1A(IIIc) specifically upregulated, from breast and pancreatic cancer to Pfeiffer syndrome and osteoarthritis (4, 9-11). A soluble version of the FGFR1A(IIIc) splice variant has shown anti-angiogenic and anti-proliferative properties in multiple cancer cell line models (11).

### References:

- Ornitz, D.M. and Itoh, N. (2015) Wiley Interdiscip. Rev. Dev. Biol. **4**:215.
- Zhang, X. *et al.* (2006) J Biol. Chem. **281**:15694.
- Ferguson, H.R. *et al.* (2021) Signaling Cells **10**:1201.
- Holzmann, K. *et al.* (2012) J Nucleic. Acids **2012**:950508.
- Xie, Y. *et al.* (2020) Sig. Transduct. Target Ther. **5**:181.
- Mossahebi-Mohammadi, M. *et al.* (2020) Front Cell Dev. Biol. **18**:79.
- Hughes, S.E. (1997) J. Histochem. Cytochem. **45**:1005.
- Delezoide, A.L. *et al.* (1998) Mech. Dev. **77**:19.
- Yamashita-Sugahara, Y. *et al.* (2016) Sci. Rep. **6**:35908.
- Teven, C.M. *et al.* (2014) Genes Dis. **1**:199.
- Babina, I. and Turner, N. (2017) Nat. Rev Cancer **17**:318.

**PRODUCT SPECIFIC NOTICES**

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.