

Monoclonal

Anti-human Siglec-7/CD328-Alexa Fluor® 488

Catalog Number: FAB1138G Lot Number: ACNJ01

100 Tests

Reagents Provided

Alexa Fluor® 488-conjugated mouse monoclonal anti-human Siglec-7/CD328: Supplied as 25 μ g of antibody in 0.5 mL saline containing up to 0.5% BSA and 0.1% sodium azide.

Clone #: 194212 Isotype: mouse IgG₂₈

Reagents Not Provided

 Flow Cytometry Staining Buffer (Catalog # FC001) or other BSA-supplemented saline buffer.

Storage


Reagents are stable for **twelve months** from the date of receipt when stored in the dark at 2° - 8° C.

Intended Use

Designed to quantitatively determine the percentage of cells bearing Siglec-7/CD328 within a population and qualitatively determine the density of Siglec-7/CD328 on cell surfaces by flow cytometry.

Product Description

This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, NS0-derived, recombinant human sialic acid binding Ig-like lectin 7 (rhSiglec-7) extracellular domain. The IgG fraction of the tissue culture supernatant was purified by Protein G affinity chromatography. The purified antibody was then conjugated to Alexa Fluor® 488 fluorochrome. Cell surface expression of Siglec-7/CD328 is determined by flow cytometry using 488 nm wavelength excitation and monitoring emitted fluorescence with a detector optimized to collect peak emissions at 515 - 545 nm.



Whole blood monocytes were stained with Alexa Fluor® 488-conjugated anti-human Siglec-7/CD328 (Catalog # FAB1138G, filled histogram) or Alexa Fluor® 488-conjugated isotype control (Catalog # IC0041G, open histogram).

Background Information

Siglec-7, also known as adhesion inhibitory receptor module-1 (AIRM-1), is a type I transmembrane glycoprotein with three extracellular Ig-like domains. It mediates sialic acid-dependent binding to cell surface ligands.

Flow Cytometry Validation

This antibody has been tested for flow cytometry using whole blood monocytes.

- Cells may be Fc-blocked with 1 μg of human IgG/10⁵ cells for 15 minutes at room temperature. Do not wash excess blocking IgG from this reaction.
- After blocking, 10 μL of conjugated antibody was added to up to 1 x 10⁶ cells and incubated for 30 minutes at room temperature.
- Unbound antibody was removed by washing the cells twice in Flow Cytometry Staining Buffer (Catalog # FC001). Note that whole blood requires a RBC lysis step at this point using Flow Cytometry Human Lyse Buffer (Catalog # FC002).
- 4. The cells were resuspended in Flow Cytometry Staining Buffer for final flow cytometric analysis. As a control for this analysis, cells in a separate tube should be treated with Alexa Fluor® 488-labeled mouse IgG_{2B} antibody. This procedure may need to be modified, depending upon the cell type and final utilization. Individual users may need to titrate to determine the optimal reagent amount for their specific use.

Warning: Contains sodium azide as a preservative - sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large volumes of water during disposal.

Legal

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose, Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.