

DESCRIPTION

Species Reactivity	Human
Specificity	Detects CD45RA in direct ELISA.
Source	Monoclonal Mouse IgG ₁ Clone # 1069433
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Whole cell transfectant expressing CD45RA Accession # P08575
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Flow Cytometry	Titration recommended for optimal concentration with starting range of 0.1-1 µg/1 million cells. Sample used for this experiment was PBMC lymphocytes.
-----------------------	--

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

This monoclonal antibody specifically binds to the 220 kDa isoform of human CD45, CD45RA. CD45RA is a single chain type I glycoprotein that is encoded by the human PTPRC gene. In humans, CD45RA is one of six isoforms of the CD45 complex. These different isoforms are the result of alternative splicing and some of the other isoforms include CD45RO, CD45RB, and CD45RC. CD45RO, CD45RB, CD45RC, and CD45RA are primarily expressed on T cells. CD45RA contains exon 4 but lacks both exons 5 and 6. CD45RO contains exons 3, 7, and 8 but not RA, RB, and RC exons. Phenotypically, CD45RA is highly expressed on naïve T cells whereas CD45RO is expressed on activated and memory T cells. Other cell types such as activated monocytes and B cells can also express CD45RO. Both CD45RA and CD45RO are used to characterize different subsets of Memory T cells.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.