

DESCRIPTION

Species Reactivity	Human
Specificity	Detects rhCDH-7 in Direct ELISA.
Source	Monoclonal Mouse IgG _{2B} Clone # 1087216
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese Hamster Ovary cell line, CHO-derived human Cadherin-7 Met1-Thr607 Accession # Q9ULB5
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cadherin-7 is an approximately 115 kDa type I transmembrane protein belonging to the Cadherin superfamily of calcium-dependent adhesion molecules. Cadherins are involved in multiple processes including embryonic development, cell migration, and maintenance of epithelial integrity (1). Human Cadherin-7 is synthesized with a 27 amino acid (aa) signal peptide and a 20 aa N-terminal propeptide. The mature cell surface-expressed protein consists of a 738 amino acid (aa) extracellular domain (ECD) that contains five Cadherin repeats, a 21 aa transmembrane segment, and a 157 aa cytoplasmic domain (2, 3). Within the ECD, human Cadherin-7 shares 99% and 97% aa sequence identity with mouse and rat Cadherin-7, respectively. Cadherin-7 interacts homotypically and heterotypically with Cadherin-14 and more weakly with Cadherins-6, -9, and -12 (3, 4). Cellular adhesion mediated by Cadherin-7 is more weak than that mediated by E- or N-Cadherin, although it can be strengthened by Fibronectin binding to Integrins on the same cell (5, 6). Cadherin-7 is localized to discrete regions of the developing nervous system. In chick and mouse, it is expressed in the basal plate of the neural tube, migrating cranial motoneurons, and migrating neural crest cells (4, 7, 8), lateral regions of the hindbrain and migrating Purkinje cell precursors (8, 9), striatum, parahippocampal areas, and somatosensory cortex of the forebrain (10 - 12), neural retina and cochlea (13, 14). Cadherin-7 interactions promote motor axon growth and inhibit axonal branching, whereas Cadherin-6B promotes branching during cranial motoneuron development (7). Cadherin-7 performs a similar function during limb bud development during which it participates in the migration and condensation of mesenchymal cells (15). Cadherin-7 is overexpressed on primary melanoma cells where it binds melanoma inhibitory activity (MIA), a secreted melanoma cell protein that promotes tumor progression (16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.