

DESCRIPTION

Species Reactivity	Human
Specificity	Detects recombinant human MAG protein in Direct ELISA.
Source	Monoclonal Mouse IgG _{2B} Clone # 1099521
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell, HEK293-derived human MAG/Siglec-4a Gly20-Pro516
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Myelin-Associated Glycoprotein (MAG), also known as Siglec-4a, is a type I transmembrane glycoprotein belonging to the Siglec family, a subgroup of the Ig superfamily (1). It is composed of an extracellular segment containing five Ig-like domains, a single transmembrane segment, and a cytoplasmic domain. Mature MAG exists as two isoforms, termed S-MAG (short) and L-MAG (long), due to alternative splicing of the cytoplasmic domain (1, 2). S-MAG has a predicted molecular weight of 67 kDa while L-MAG has a predicted molecular weight of 71 kDa (1, 2). Additionally, proteolytic cleavage of the extracellular domain produces a soluble MAG (3). Within shared regions in the extracellular domain, human MAG shares 95% aa sequence identity with mouse and rat MAG. MAG functions as an adhesion molecule during neural development. It preferentially binds to α-2,3-linked sialic acid terminal structures found on cell surface molecules (1, 4, 5). MAG is selectively expressed by myelinating oligodendrocytes and Schwann cells and plays an important role in axon-myelin stability (1, 4). Specifically, L-MAG is involved in myelination in the central nervous system (CNS) while S-MAG is the predominate isoform expressed during myelination in the peripheral nervous system (1). MAG is also reported to regulate the axon cytoskeleton and support the distribution of axon molecules at the nodes of Ranvier (1, 4). In addition, it has been identified as a major inhibitor of neurite outgrowth (1, 4, 6). However, MAG has also been reported to protect neurons from excitotoxicity (1, 7). MAG is believed to utilize the gangliosides GD1a and GT1b, the Nogo receptors NgR1 and NgR2/NGRH1, Integrin β1/CD29, and PIR-B to mediate its effects (1, 4, 5, 8, 9). Soluble MAG, which is released from myelin in large quantities, has been identified in normal human tissues and in tissues from patients with neurological disorders (3). It is believed that this soluble MAG might contribute to the lack of CNS neuron regeneration after injury (3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.