

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-20 R α in direct ELISAs and Western blots. In Western blots, approximately 5-15% cross-reactivity with recombinant human (rh) IFN- γ R2, rhIL-10 R, rhIL-10 R β , rhIL-20 R β , and rhIL-22 BP is observed and no cross-reactivity with rhIFN- γ RI, recombinant mouse IL-20 R α , or rhIL-22 R is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 173714
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-20 R α Val30-Lys250 Accession # Q9UHF4
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 μ g/10 ⁶ cells	Human IL-20 R α transfected Baf/3 cells

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

IL-20 receptor alpha (IL-20 R α), also named IL-20 R1, CRF2-8, and ZCYTOR7, belongs to the class II cytokine receptor family, which includes 12 members. These receptors are characterized by the patterns of conserved amino acid (aa) residues in their extracellular domains, which are composed of tandem fibronectin type III domains (1). Class II cytokine receptors form heterodimeric signaling receptor complexes that mediate class II cytokine signals. Subunits of the different receptor complexes are shared and serve multiple functions (1).

The gene for human IL-20 R α is mapped to chromosome 6 and encodes a 553 aa glycoprotein with a 29 aa signal peptide, a 221 aa extracellular domain, a 24 aa transmembrane region and a 279 aa intracellular domain (2). IL-20 R α is widely expressed and is detected at high levels in multiple tissues including skin, testis, heart, placenta, salivary gland and prostate gland (1). The expression of IL-20 R α , together with that of IL-20 R β , is upregulated in psoriatic skin lesions on keratinocytes, immune cells, and endothelial cells (1, 2).

IL-20 R α heterodimerizes with IL-20 R β to form the functional receptor that mediates IL-19, IL-20 and IL-24 signals (3, 4). IL-20 R α also heterodimerizes with IL-10 R β to form the functional receptor complex for IL-26 (5). Binding of these IL-10 family class II cytokines to their functional receptors induces activation of the JAK-STAT signal transduction pathway. At low ligand concentrations, STAT3 has been shown to be the predominant STAT proteins activated through either complexes (3-5).

References:

1. Kotenko, S.V. (2003) *Cytokine & Growth Factor Reviews* **13**:223.
2. Xie, M.H. *et al.* (2000) *J. Biol. Chem.* **275**:31335.
3. Dumoutier, L. *et al.* (2001) *J. Immunol.* **167**:3534.
4. Parrish-Novak, J. *et al.* (2002) *J. Biol. Chem.* **277**:47517s.
5. Sheikh, F. *et al.* (2004) *J. Immunol.* **172**:2006.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.