

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-18 R β /IL-1 R7 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 1% cross-reactivity with recombinant mouse IL-18 R β , recombinant human (rh) IL-18 R, rhIL-1 RI, rhIL-1 RII, rhIL-1 RAcP, and rhIL-1 Rrp2
Source	Monoclonal Mouse IgG ₁ Clone # 132016
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-18 R β /IL-1 R7 Met1-Arg356 Accession # O95256
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-18, originally described as an interferon- γ inducing factor (IGIF), is a member of the IL-1 family of cytokines that has multiple immunoregulatory functions. It has potent IFN- γ inducing activities and plays a key role in the activation of T helper type 1 (Th1) responses. The functional IL-18 receptor complex consists of two components, the IL-18 R α (IL-1 R5) and IL-18 R β (also termed IL-1 R7 and AcPL) subunits. Both subunits are members of the IL-1 receptor superfamily. Although IL-18 R α by itself binds IL-18 with low-affinity and IL-18 R β does not bind IL-18 *in vitro*, co-expression of IL-18 R α and IL-18 R β is required for high-affinity binding and IL-18 responsiveness. Human IL-18 R β cDNA encodes a 599 amino acid (aa) residue precursor type I membrane protein with a 14 aa signal peptide, a 342 aa extracellular region containing three immunoglobulin-like domains, a single transmembrane domain and a 222 aa cytoplasmic domain. Human and mouse IL-18 R β share 65% aa sequence identity. The expression of IL-18 R β parallels that of IL-18 R α and is detected in numerous tissues including lung, spleen, leukocytes and colon.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.