

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-18 BPα in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse (rm) IL-18 BPc or rmlIL-18 BPD is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 136007
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-18 BPα Thr29-Gly192 Accession # AAD17190
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 18 binding protein (IL-18 BP or IL-18 BPα) is a secreted glycoprotein, which functions as an IL-18 antagonist by binding to IL-18 and blocking its biological activity. IL-18 BPα bears no amino acid sequence homology to the membrane-associated IL-18 and IL-1 receptor proteins. The gene for human IL-18 BPα has been localized to chromosome 11q13. It encodes for at least four isoforms by alternative splicing. The IL-18 BP isoforms a and c each contain one immunoglobulin (Ig)-like C2-type domain while isoforms b and d lack a complete Ig domain. The complete Ig domain has been shown to be essential to the binding and neutralizing properties of the binding proteins. Two isoforms of mouse IL18 BP (c and d) containing the complete Ig domain have also been isolated and shown to neutralize IL-18 bioactivity. Human and mouse IL-18 BPα share approximately 61% amino acid sequence identity. Several poxviruses also encode proteins with sequence similarity to the human and mouse IL-18 BPα. Viral IL-18 BPs have been shown to bind and inhibit IL-18 responses and may be involved in modulating host immune responses. The expression of IL-18 BP is markedly upregulated by IFN-γ, suggesting that IL-18 activity is modulated by a negative feedback mechanism mediated by IL-18 BP.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.