

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-20 in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 380605
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse IL-20 Leu25-Leu176 Accession # Q9JKV9.1
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Mouse Interleukin 20 (IL-20) was identified by searching sequence databases for translated sequences with a signal sequence and amphipathic helices found in helical cytokines. Based on the human molecule, mouse IL-20 was discovered in a skin library. Mouse IL-20 is synthesized as a 176 amino acid (aa) precursor that contains a 24 aa signal sequence and a 152 aa mature segment. There are no N-linked glycosylation sites and it is doubtful that the native molecule is glycosylated. Although IL-20 is a distant member of the IL-10 family, it functions as a monomer. IL-20 shares less than 40% aa sequence identity with other IL-10 family members. Mouse and human IL-20 are 77% aa identical in the mature segment. IL-20 production has been found in skin and trachea. In particular, activated keratinocytes and, possibly, monocytes are reported to express IL-20. There are two heterodimeric receptor complexes for IL-20. The first complex is composed of IL-20 R α and IL-20 R β . The second complex is composed of IL-22 R and IL-20 R β . Whereas the IL-22 R/IL-20 R β complex is shared with IL-24/mda-7, the IL-20 R α /IL-20 R β complex is shared with both IL-19 and IL-24. Little is known about the function of IL-20. It is reported to induce the proliferation of multipotential hematopoietic progenitor cells, direct the differentiation and expansion of keratinocytes, and promote the release of pro-inflammatory mediators in keratinocytes and other IL-20 receptor expressing cells (1-6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.