

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FGF-3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) FGF-4, -5, -6, -7, -9, -10, -11, -12, -16, -17, -18, -19, -20, -21, -23, rhFGF acidic, rhFGF ba
Source	Monoclonal Mouse IgG _{2A} Clone # 254625
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human FGF-3 Asp28-Arg212 Accession # P11487
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fibroblast Growth Factor 3 (FGF-3) belongs to the large FGF family which has at least 23 members (1, 2). All FGF family members are heparin-binding growth factors with a core 120 amino acid (aa) FGF domain that allows for a common tertiary structure. FGFs are expressed during embryonic development and in restricted adult tissues. They act on cells of mesodermal and neuroectodermal origin to regulate diverse physiologic functions including angiogenesis, cell growth, pattern formation, embryonic development, metabolic regulation, cell migration, neurotrophic effects and tissue repair (3, 4). Signaling receptors for FGFs are type I transmembrane receptor tyrosine kinases belonging to the Ig superfamily. Four distinct but related classes of FGF receptors, FGF R1, 2, 3, and 4, exist. Through alternative splicing, multiple isoforms for FGF R1, 2 and 3, with distinct ligand recognition profiles, are also generated (4).

The *FGF-3* gene, originally designated *int-2*, was first identified as a proto-oncogene activated in mouse mammary tumors by proviral integration (2). Amplification of this gene has also been found frequently in human tumors. Human FGF-3 cDNA predicts a 239 aa precursor protein with a 17 aa signal peptide and a 222 aa secreted mature protein with one potential N-linked glycosylation site (1). Human and mouse FGF-3 share 88% aa sequence identity. The *Xenopus* and mammalian secreted FGF-3 are processed proteolytically at both the N- and C-terminus (5). FGF-3 binds with high-affinity to the IIIb isoforms of FGF R1 and FGF R2. FGF-3 also binds the IIIc isoform of FGF R2, but with lower affinity (6). FGF-3 has been implicated in the induction of inner ear development (7). Studies have suggested that FGF-3 and FGF-8 function synergistically in otic placode induction and during neuronal development to regulate dorsoventral axis formation (8 - 10). During development, the activities of FGF-3 and FGF-8 are regulated negatively by the sprouty family proteins and by Sef (similar expression to *fgf* genes), a transmembrane protein that shares intracellular sequence similarities with the IL-17 receptor (10).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.