

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Nogo Receptor/NgR in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse Nogo Receptor/NgR is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 188428
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human Nogo Receptor/NgR Cys27-Ser447 (predicted) Accession # Q9BZR6
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Nogo Receptor (NgR), also named reticulon 4 receptor, is a glycosylphosphoinositol (GPI)-anchored protein that belongs to the family of leucine-rich repeat (LRR) proteins (1). It is expressed predominantly in the central nervous systems in neurons and their axons. NgR plays an essential role in mediating axon growth inhibition induced by the structurally distinct myelin-derived proteins Nogo, myelin-associated glycoprotein (MAG), and myelin oligodendrocyte glycoprotein (Omgp) (2, 3). Human NgR cDNA encodes a 473 amino acid (aa) precursor with a 26 aa putative signal peptide, an LRR-type N-terminal region, eight LRR repeats, a cysteine-rich LRR-type C-terminal region, a GPI linkage domain and a 26 aa C-terminal propeptide that is removed in the mature form (1). All of the LRR domains within NgR are required for ligand binding and receptor oligomerization (4). NgR mediates its inhibitory actions by interacting with the p75 neurotrophin receptor (p75^{NTR}), a tumor necrosis factor receptor superfamily (TNFRSF) member also known for modulating the activities of the Trk family of receptor tyrosine kinases, and for inducing apoptosis in neurons and oligodendrocytes (5). Upon ligand binding, NgR binds to and activates the p75^{NTR}. The activated p75^{NTR} then sequesters the Rho guanine dissociation inhibitor (Rho-GDI) away from Rho and allows Rho to change into the active GTP-bound state which can interact with signaling proteins to suppress axonal growth and regeneration (4). The truncated extracellular domain of NgR has been shown to bind the myelin-derived inhibitors and block inhibition of axon growth by myelin (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.