

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse Sca-1/Ly6 in direct ELISAs.
Source	Monoclonal Rat IgG _{2A} Clone # 177228
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Sca-1/Ly6 C-terminally truncated Ly-6E allele Leu27-Gly119 Accession # CAA28351
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Stem Cell Antigen-1 (Sca-1) is encoded by the strain-specific *Ly-6 E/A* allelic gene. Its expression on multipotent hematopoietic stem cells (HSC) has been used as a marker of HSC in mice of both *Ly-6* haplotypes (2, 3). This antibody is frequently used in combination with lineage depletion antibodies to identify and isolate murine HSC. Sca-1-positive HSC can be found in the adult bone marrow, fetal liver and mobilized peripheral blood and spleen in the adult animal (2-7). However, Sca-1 has also been discovered in several non-hematopoietic tissues (1) and can be used to enrich progenitor cell populations other than HSC (8). It is suggested that Sca-1 could be involved in regulating both B and T cell activation (9-12).

References:

1. Van de Rijn, M. *et al.* (1989) Proc. Natl. Acad. Sci. USA **86**:4634.
2. Spangrude, G.I. *et al.* (1988) Science **241**:58.
3. Spangrude, G.I. *et al.* (1993) Blood **82**:3327.
4. Morrison, S.J. *et al.* (1995) Proc. Natl. Acad. Sci. USA **92**:10302.
5. Kawamoto, H. *et al.* (1997) Int. Immunol. **9**:1011.
6. Yamamoto, Y. *et al.* (1996) Blood **88**:445.
7. Morrison, S.J. *et al.* (1997) Proc. Natl. Acad. Sci. USA **94**:1908.
8. Welm, B.E. *et al.* (2002) Dev. Biol. **245**:42.
9. Codias, E.K. *et al.* (1990) J. Immunol. **144**:2197.
10. Malek, T.R. *et al.* (1986) J. Exp. Med. **164**:709.
11. Codias, E.K. *et al.* (1990) J. Immunol. **145**:1407.
12. Flood, P.M. *et al.* (1990) J. Exp. Med. **172**:115.

