

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse OX40 Ligand/TNFSF4 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) APRIL, rhGITR Ligand, rhLIGHT, rhLTα1/β2, rhLTα2/β1, recombinant cotton rat TNF-α, recombinant mo
Source	Monoclonal Rat IgG _{2A} Clone # 182601
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse OX40 Ligand/TNFSF4 Gln49-Leu198 Accession # P43488
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

OX40 Ligand (OX40L), also known as gp34, is a type II transmembrane glycoprotein belonging to the TNF superfamily. Murine OX40L cDNA encodes a 198 amino acid (aa) protein comprised of a 28 aa N-terminal cytoplasmic domain, a 20 aa transmembrane segment, and a 150 aa C-terminal extracellular domain (1). Human and mouse OX40L share 46% sequence identity at the amino acid level (1). OX40L is expressed on activated antigen presenting cells such as B cells, macrophages, dendritic cells, and on endothelial cells at the site of inflammation. The receptor for OX40L is OX40 (CD134) which is expressed predominantly on activated CD4⁺ T cells. Expression of OX40 is transient following engagement of T cell receptors (2). Ligation of OX40L by OX40 stimulates proliferation and differentiation of activated B cells, and increases immunoglobulin secretion (3, 4). The expression of OX40L on B cells is upregulated by CD40 ligation (3). Engagement of the OX40-OX40L system has co-stimulatory effects on T cells by stimulating the production of cytokines by T helper cells and increasing the survival of memory T cells (2, 5). Blocking of the OX40-OX40L interaction *in vitro* inhibits co-stimulation resulting in decreased T cell proliferation and adhesion of T cells to endothelial cells. Inhibition of the OX40-OX40L interaction in disease models has beneficial effects in acute graft-versus-host disease, inflammatory bowel disease and decreases the development of collagen-induced arthritis and experimental leishmaniasis (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.