

DESCRIPTION

Species Reactivity	SARS-CoV-2
Specificity	Detects SARS-CoV-2 Spike RBD in direct ELISAs.
Source	Recombinant Monoclonal Rabbit IgG Clone # 2820D
Purification	Protein A or G purified
Immunogen	Human embryonic kidney cell HEK293-derived SARS-CoV-2 Spike RBD Protein Ala319-Phe541 Accession # YP_009724390.1
Conjugate	Alexa Fluor Plus 488 Excitation Wavelength: 493 nm Emission Wavelength: 518 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

DATA

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SARS-CoV-2, which causes the global pandemic coronavirus disease 2019 (Covid-19), belongs to a family of viruses known as coronaviruses that are commonly comprised of four structural proteins: Spike protein(S), Envelope protein (E), Membrane protein (M), and Nucleocapsid protein (N) (1). SARS-CoV-2 Spike Protein (S Protein) is a glycoprotein that mediates membrane fusion and viral entry. The S protein is homotrimeric, with each ~180-kDa monomer consisting of two subunits, S1 and S2 (2). In SARS-CoV-2, as with most coronaviruses, proteolytic cleavage of the S protein into two distinct peptides, S1 and S2 subunits, is required for activation. The S1 subunit is focused on attachment of the protein to the host receptor while the S2 subunit is involved with cell fusion (3-5). Based on structural biology studies, the receptor binding domain (RBD), located in the C-terminal region of S1, can be oriented either in the up/standing or down/lying state (6). The standing state is associated with higher pathogenicity and both SARS-CoV-1 and MERS can access this state due to the flexibility in their respective RBDs. A similar two-state structure and flexibility is found in the SARS-CoV-2 RBD (7). Based on amino acid (aa) sequence homology, the SARS-CoV-2 S1 subunit RBD has 73% identity with the RBD of the SARS-CoV-1 S1 RBD, but only 22% homology with the MERS S1 RBD. The low aa sequence homology is consistent with the finding that SARS and MERS bind different cellular receptors (8). The S Protein of the SARS-CoV-2 virus, like the SARS-CoV-1 counterpart, binds Angiotensin-Converting Enzyme 2 (ACE2), but with much higher affinity and faster binding kinetics (9). Before binding to the ACE2 receptor, structural analysis of the S1 trimer shows that only one of the three RBD domains in the trimeric structure is in the "up" conformation. This is an unstable and transient state that passes between trimeric subunits but is nevertheless an exposed state to be targeted for neutralizing antibody therapy (10). Polyclonal antibodies to the RBD of the SARS-CoV-2 protein have been shown to inhibit interaction with the ACE2 receptor, confirming RBD as an attractive target for vaccinations or antiviral therapy (11). There is also promising work showing that the RBD may be used to detect presence of neutralizing antibodies present in a patient's bloodstream, consistent with developed immunity after exposure to the SARS-CoV-2 virus (12). Lastly, it has been demonstrated the S Protein can invade host cells through the CD147/EMMPRIN receptor and mediate membrane fusion (13, 14).

References:

1. Wu, F. *et al.* (2020) Nature **579**:265.
2. Tortorici, M.A. and D. Veasler (2019). Adv. Virus Res. **105**:93.
3. Bosch, B.J. *et al.* (2003) J. Virol. **77**:8801.
4. Belouzard, S. *et al.* (2009) Proc. Natl. Acad. Sci. **106**:5871.
5. Millet, J.K. and G. R. Whittaker (2015) Virus Res. **202**:120.
6. Yuan, Y. *et al.* (2017) Nat. Commun. **8**:15092.
7. Walls, A.C. *et al.* (2010) Cell **180**:281.
8. Jiang, S. *et al.* (2020) Trends. Immunol. <https://doi.org/10.1016/j.it.2020.03.007>.
9. Ortega, J.T. *et al.* (2020) EXCLI J. **19**:410.
10. Wrapp, D. *et al.* (2020) Science **367**:1260.
11. Tai, W. *et al.* (2020) Cell. Mol. Immunol. <https://doi.org/10.1016/j.it.2020.03.007>.
12. Okba, N. M. A. *et al.* (2020). Emerg. Infect. Dis. <https://doi.org/10.3201/eid2607.200841>.
13. Wang, X. *et al.* (2020) <https://doi.org/10.1038/s41423-020-0424-9>.
14. Wang, K. *et al.* (2020) bioRxiv <https://www.biorxiv.org/content/10.1101/2020.03.14.988345v1>.

PRODUCT SPECIFIC NOTICES

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.