

#### DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human PDGF R $\beta$ in Western blots. Specificity has been confirmed in binding studies using several different cell lines (1, 2) and by its ability to immunoprecipitate PDGF receptor $\beta$ -subunit complexed with <sup>125</sup> I-PDGF-BB (3). It does not recognize the PDGF receptor $\alpha$ -subunit. Detects the PDGF receptor $\beta$ -subunit of human and primate species (monkey and baboon) but not the rat or mouse receptors. Its ability to bind to receptors from other species has not been tested.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # PR7212
<b>Purification</b>	Protein A or G purified from ascites
<b>Immunogen</b>	Human skin fibroblast membrane extracts
<b>Conjugate</b>	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	0.25-1 $\mu$ g/10 <sup>6</sup> cells	BUD-8 human fibroblast cell line

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> ● 12 months from date of receipt, 2 to 8 °C as supplied.

#### BACKGROUND

PDGF is a major serum mitogen that can exist as a homo or hetero-dimeric protein consisting of disulfide-linked PDGF-A and PDGF-B chains. The PDGF-AA, PDGF-BB and PDGF-AB isoforms have been shown to bind to two distinct cell surface PDGF receptors with different affinities. Where as PDGF R $\alpha$  binds all three PDGF isoforms with high affinity, PDGF R $\beta$  binds PDGF-BB only with high-affinity. Both PDGF R $\alpha$  and PDGF R $\beta$  are members of the class III subfamily of receptor tyrosine kinases (RTK) that also includes the receptors for M-CSF, SCF and Flt3 ligand. All class III RTKs are characterized by the presence of five immuno-globulinlike domains in their extracellular region and a split kinase domain in their intracellular region. PDGF binding induces receptor homo-and hetero-dimerization and signal transduction. The expression of the  $\alpha$  and  $\beta$  receptors is independently regulated in various cell types. Recombinant soluble PDGF R $\beta$  binds PDGF with high affinity and is potent PDGF antagonist (4).

#### References:

- Hart *et al.* (1987) J. Biol. Chem. **262**:10780.
- Gronwald *et al.* (1988) Proc. Natl. Acad. Sci. **85**:3435.
- Seifert *et al.* (1989) J. Biol. Chem. **264**:8771.
- Heldin, C.H. and L. Claesson-Welsh (1994) in *Guidebook to Cytokines and Their Receptors*, Nicola, N.A. ed. Oxford University Press, New York, p. 202.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.