

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Jagged 1 in direct ELISAs. In direct ELISAs, approximately 50% cross-reactivity with recombinant human Jagged 2 and recombinant rat Jagged 1 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 188323
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0-derived recombinant human Jagged 1 Ser32-Asp296 Accession # P78504
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Jagged 1 is a 180 kDa type I transmembrane glycoprotein and member of the Delta-Serrate-Lag-2 (DSL) family of ligands that activate LIN12/Notch proteins. Human Jagged 1 is synthesized as a 1218 amino acid (aa) precursor that contains a 33 aa signal sequence, a 1034 aa extracellular domain (ECD), a 26 aa transmembrane segment, and a 125 aa cytoplasmic region. The ECD contains a DSL domain (aa 185-229), a cysteine-rich region, 15 EGF-like repeats, of which many bind calcium, and nine potential sites for N-linked glycosylation. Mature human Jagged 1 shares 97% and 96% aa identity with mature mouse and rat Jagged 1, respectively. Jagged 1 is widely expressed in adult and fetal tissues. Jagged-Notch signaling specifies cell fate, regulates pattern formation, defines boundaries between different cell types, and modulates cell proliferation and differentiation, especially during hematopoiesis, myogenesis, neurogenesis, and development of vasculature (1-8). Mutations in human Jagged 1 are the cause of Alagille syndrome, an autosomal-dominant disorder characterized by intrahepatic cholestasis and abnormalities of heart, eye, vertebrae, as well as characteristic facial appearance (9, 10).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.