

DESCRIPTION

Species Reactivity	Chicken
Specificity	Detects chicken Netrin-2 in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 74506
Purification	Protein A or G purified from ascites
Immunogen	Mouse myeloma cell line NS0-derived recombinant chicken Netrin-2 Ala16-Pro581 Accession # Q90923
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Netrins are laminin-like axon guidance molecules that are conserved among *C. elegans*, *Drosophila*, and vertebrates. The chicken Netrin-2 cDNA encodes a precursor protein that has structural similarity to the N-terminus of B2 or the γ-chain of laminin. The mature Netrin-2 protein has 581 amino acid (aa) which contains a 245 aa domain VI, a 169 aa domain V with three cysteine/glycine rich EGF repeats and a unique 152 aa domain C that enriches in basic aa residues and contains an RGD motif. Chicken Netrin-2 shares 72% aa identity with chicken Netrin-1, 51% aa identity with *C. elegans* UNC-6 protein and 57% aa identity with human NTN2L (Netrin-2-like). Although only two chicken netrins are known, more mammalian netrins are known. Netrin-2 is expressed in the chick brain and spinal cord from E6 through to the adult. In addition, expression of Netrin-2 was observed at various stages in several mesodermally and endodermally derived tissues, including lung, gut, ovary, testes and spleen. Netrins act to both attract and repel the growing axons of a broad range of neuronal cell types during development. These actions are mediated by specific receptor complexes containing either the DCC (colorectal carcinoma) or neogenin, in the case of attractant, or UNC-5-related proteins, in the case of repellent.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.