

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Follistatin-related Gene Protein/FLRG in ELISAs and Western blots. In sandwich immunoassays and Western blots, no cross-reactivity with recombinant mouse FLRG is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 206714
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Follistatin-related Gene Protein/FLRG Met27-Val263 Accession # O95633
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair) Optimal dilution of this antibody should be experimentally determined.

ELISA Detection (Matched Antibody Pair) Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Follistatin-related gene protein (FLRG), also known as follistatin-like 3 (FSTL3) is a glycoprotein belonging to the follistatin-module protein family. Human FLRG cDNA encodes a 263 amino acid (aa) residue protein with a putative 26 aa signal peptide, an N-terminal domain, two cysteine-rich follistatin-like domains (FS) and a C-terminal acidic domain. Compared to follistatin, FLRG lacks the third FS domain found in follistatin. In addition, FLRG also lacks the heparin-binding domain found within the first amino-terminal FS domain of follistatin. Mouse and human FLRG share approximately 83% aa sequence homology. Like follistatin, FLRG has been shown to bind and inhibit the activities of TGF-β family ligands including activin, BMP-2, -6, -7 and GDF-8/myostatin. While both FLRG and follistatin are located in a wide and overlapping range of adult and fetal tissue, their sites of peak expression differ: FLRG most highly in heart, lung, kidney, placenta and testis, while follistatin is highest in ovary and pituitary. The expression of FLRG is upregulated by TGF-β and activin signaling through Smad proteins. Although FLRG is a secreted protein in many cell types, it has also been localized to the nuclear compartment in HeLa, 293 and CHO cells (1-5).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.