

## DESCRIPTION


<b>Species Reactivity</b>	Human/Mouse/Bovine
<b>Specificity</b>	Detects human, mouse, and bovine Integrin $\alpha 6$ /CD49f. Recognizes an epitope in the extracellular domain of the Integrin $\alpha 6$ subunit.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # GoH3
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse mammary tumor cells
<b>Conjugate</b>	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

The Integrin  $\alpha 6$  subunit, also known as CD49f and VLA-6  $\alpha$  subunit, forms a heterodimer with Integrin  $\beta 1$  (CD29) or  $\beta 4$  (CD104) subunits and recognizes laminin as a ligand.