

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human NKG2D/CD314 in direct ELISA.
Source	Recombinant Monoclonal Rabbit IgG Clone # 2855A
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Mouse myeloma cell line, NS0-derived human NKG2D/CD314 Phe78-Val216 Accession # P26718
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS	
Please Note: Optimal dilutions should be	determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.
Flow Cytometry	Titration recommended for optimal concentration with starting range of 0.1-1 μg/1 million cells. Sample used for this experiment was NK cells activated and expanded with CD56 costain for 10 days.

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze.	
	 12 months from date of receipt, 2 to 8 °C as supplied. 	

BACKGROUND

NKG2D is a type II transmembrane glycoprotein having an extracellular lectin-like domain. This domain lacks the recognizable calcium-binding sites found in true C-

type lectins and binds protein rather than carbohydrate ligands. Human NKG2D is expressed on CD8⁺ αβ T cells, γδ T cells, NK cells and NKT cells. In mouse systems NKG2D also occurs on macrophages. Human ligands for NKG2D include MICA, MICB, and ULBP1, 2, and 3. Expression of NKG2D ligands occurs in epithelial cells, tumor cells and under conditions of stress or infection. NKG2D exists as a disulfide-linked homodimer that delivers an activating signal upon ligand binding. Signaling requires association with an adapter protein. Alternative splicing of the NKG2D mRNA results in isoforms with different cytoplasmic domains that can associate either with DAP12 to deliver a true activating signal or with DAP10 resulting in a costimulatory signal. NKG2D has been implicated in anti-tumor surveillance and the immune response against viral infection.

References:

- 1. Li, P. *et al*. (2001) Nature Immunol. 2:443.
- 2. Steinle, A. et al. (2001) Immunogenetics 53:279.
- 3. Cosman, D. et al. (2001) Immunity 14:123.
- 4. Cerwenka, A. and L. Lanier (2001) Immunol. Rev. 181:158.
- 5. Wu, J. et al. (1999) Science 285:730.
- 6. Diefenbach, A. et al. (2002) Nature Immunol. 3:1142.
- 7. Gilfillan, S. et al. (2002) Nature Immunol. 3:1150.
- 8. Groh, V. et al. (2001) Nature Immunol. 2:255.
- 9. Cerwenka, A. et al. (2001) Proc. Natl. Acad. Sci. USA 98:11521.
- 10. Diefenbach, A. et al. (2001) Nature 413:165.
- 11. NKG2D and its Ligands (2002) www.rndsystems.com

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 1/23/2023 Page 1 of 1

biotechne^{GIODAL DIO-TECHNE.COT} USA TEL 800 343 74 Europe I Middle East

Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449