

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Cryptic in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse Cryptic is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 219708
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Cryptic Tyr26-Gly169 Accession # P0CG36
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cryptic, also known as CFC-1, was originally identified as a secreted protein that is specifically expressed during mesoderm differentiation (1). Cryptic, along with Cripto, *Xenopus* FRL-1 and zebrafish OEP (one-eyed pinhead) define the epidermal growth factor-CriptoFRL1Cryptic (EGF-CFC) family of signaling proteins that function in cancer and various developmental processes (2, 3). Overall sequence identity between members of the family is low, but they do share distinct domains: a variant EGF-like motif, a novel conserved cysteine-rich domain (called CFC domain), and a C-terminal hydrophobic region (2). Most EGF-CFC members have a glycosyl-phosphatidylinositol (GPI) anchoring site at the C-terminus and exist as extracellular membrane-anchored proteins. However, naturally-occurring soluble isoforms also exist. Human Cryptic shares 55% and 25% amino acid identity with mouse Cryptic and human Cripto, respectively. Despite weak conservation in amino acid identity, EGF-CFC family members appear to function similarly in assays for phenotypic rescue of zebrafish *oep* mutants (2). Cryptic is expressed during gastrulation in the mesoderm and later in the neuroectoderm, marking the prospective floor plate of the neural tube (1). Genetic evidence from mice and humans points to a role for Cryptic in determining left-right asymmetry. Mutations in the *cryptic* gene result in a spectrum of heart, lung and spleen defects, all representing left-right laterality defects (4, 5). These phenotypes resemble some Nodal mutant alleles suggesting that Cryptic, like Cripto, acts as an essential cofactor for Nodal signaling (1, 3). Studies have shown that other TGF-β superfamily members involved in mesoderm induction and left-right patterning, Vg1 and GDF-1, also require EGF-CFC cofactors. Cryptic binds to GDF-1 leading to an Act RIIb-ALK4-Cryptic-GDF-1 complex for signaling (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.