

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human IL-13 R α 1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse IL-13 R α 1, recombinant human (rh) IL-13 R α 2, rhIL-4 R, rhIL-5 R β , or rhIL-9 R is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 419718
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-13 R α 1 Ala27-Thr343 Accession # P78552
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Two type 1 membrane proteins belonging to the hemopoietin receptor family have been cloned and shown to bind IL-13 with differing affinities. The lower affinity IL-13 binding protein, previously designated IL-13 R α , IL-13 R α or NR4, is now referred to as IL-13 R α 1. The high-affinity IL-13 binding protein, previously also designated IL-13 R or IL-13 R α ', is now referred to as IL-13 R α 2.

The human IL-13 R α 1 was originally cloned based on sequence homology to the mouse IL-13 R α 1. The IL-13 R α 1 cDNA encodes a 427 amino acid (aa) precursor protein with a putative 21 aa signal peptide, a 324 aa extracellular domain, a 23 aa transmembrane region and a 59 aa cytoplasmic tail. Human and mouse IL-13 R α 1 share 76% aa sequence identity. The extracellular domain of IL-13 R α 1 is also closely related to that of IL-13 R α 2. IL-13 R α 1 has been shown to combine with the IL-4 R α to form a high-affinity receptor complex capable of transducing an IL-13-dependent proliferative signal. The role of IL-13 R α 2 in IL-13 signaling remains to be elucidated.

References:

1. Caput, D. *et al.* (1996) *J. Biol. Chem.* **271**:16921.
2. Donaldson, D.D. *et al.* (1998) *J. Immunol.* **161**:2317.
3. Aman, M.J. *et al.* (1996) *J. Biol. Chem.* **271**:29265.
4. Hilton, D.J. *et al.* (1996) *Proc. Natl. Acad. Sci. USA* **93**:497.
5. Zhang, J.G. *et al.* (1997) *J. Biol. Chem.* **272**:9474.