

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Furin in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 222728
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Furin Asp108-Glu715 Accession # NP_002560
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Furin is a member of the proprotein convertase (PC) family, which belongs to the subtilisin superfamily of serine protease (1-3). As a cellular protease, Furin processes a variety of proproteins in secretory pathway compartments by cleaving after Arg-Xaa-Lys/Arg-Arg-like motifs, which usually reside at the end of the pro regions of these proproteins. Examples of the proprotein substrates are growth factors and receptors, extracellular matrix proteins, and other proteases. Furin has an essential role in embryogenesis and homeostasis and is implicated in various pathologies such as cancer, neurodegenerative diseases and anthrax. It is synthesized as a 794 amino acid type I transmembrane protein precursor with a signal peptide (residues 1-24), a pro region (residues 25-107), which play a crucial role in the folding, activation and transport of Furin, and a mature chain (residues 108-794) (1-3). The mature chain consists of the subtilisin-like catalytic domain, a P domain, which is essential for enzyme activity and the modulation of pH and calcium requirements, and a cytoplasmic domain, which controls the localization and sorting of Furin in the *trans*-Golgi network/endosomal system. The purified recombinant human Furin (residues 108-715) corresponds to the mature enzyme terminated before the transmembrane domain.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.