

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat L-Selectin/CD62L in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody does not cross-react with recombinant mouse (rm) L-Selectin or rhL-Selectin.
Source	Monoclonal Mouse IgG ₁ Clone # 196021
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat L-Selectin/CD62L Trp39-Asn332 Accession # P30836
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

L-Selectin (also known as Leukocyte Selectin, LAM-1, LECAM-1, LECCAM-1, TQ1, Leu-8, MEL-14 antigen, DREG, lymph node homing receptor, CD62L) is a member of the Selectin family of cell surface molecules which include E-Selectin and P-Selectin. All Selectins have an extracellular domain composed of an amino-terminal calcium-dependent lectin domain, an epidermal growth factor (EGF)-like domain, two to nine short consensus repeat (SCR) units, a transmembrane domain, and a cytoplasmic tail. L-Selectin expression is limited to hematopoietic cells, with most leukocytes expressing L-Selectin at some stage of differentiation. The majority of myeloid cells, B cells, and virgin T cells express L-Selectin, while only a sub-population of memory T cells and NK cells express L-Selectin. Lymphocytes and neutrophils exhibit a reversible loss of L-Selectin after cellular activation that results from endoproteolytic release of the extracellular portion of receptor from the cell surface. Cleavage of L-Selectin from the cell surface results in a high circulating level of functionally active soluble L-Selectin. All selectins bind sialylated and fucosylated oligosaccharides that are linked to glycoproteins and glycolipids. L-Selectin specifically binds to at least three different heavily glycosylated mucin-like proteins: GlyCAM-1, CD34, and MAdCAM-1. Multiple studies indicated that L-Selectin, P-Selectin E-Selectin collaborate to mediate the initial binding of leukocytes to endothelium at sites of tissue injury and inflammation, producing the characteristic "rolling" of leukocytes along the endothelium. L-Selectin knockout mice have a 70% decrease in rolling leukocytes in exposed mesentery and have impaired neutrophil and monocyte migration into areas of inflammation. Additionally, L-Selectin knockout mice have relatively few lymphocytes present in peripheral lymph nodes and Peyer's patches. Short-term in vivo homing experiments in L-Selectin deficient mice demonstrate that L-Selectin is involved in lymphocyte homing to Peyer's patches and mesenteric lymph nodes in the gut. Rat and human L-Selectin share 77% amino acid sequence homology. Rat and mouse L-Selectin share 83% amino acid sequence homology (1, 2).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.