

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Soggy-1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody does not cross-react with recombinant mouse Soggy-1.
Source	Monoclonal Mouse IgG ₁ Clone # 202509
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Soggy-1 Ala31-Leu242 Accession # Q9UK85.1
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Soggy-1, also known as DkkL1 (Dickkopf-like 1), is a secreted protein that is related to the Dickkopf family of Wnt antagonists. Dkk-1, -2, -3, and -4 each have two cysteine-rich domains separated by a linker. Soggy-1 does not contain cysteine-rich domains but bears some sequence homology with the N-terminal regions of Dkk proteins (1, 2). The human Soggy-1 cDNA encodes a 242 amino acid (aa) precursor that includes a 30 aa signal sequence (2). Human Soggy-1 shares approximately 65% aa sequence identity with mouse and rat Soggy-1 and 15 - 20% aa sequence identity with human Dkk-1, -2, -3, and -4. Mouse Soggy-1 is expressed at various sites in the embryo but in the adult is primarily found in the testes (3 - 5). Soggy-1 transcription is regulated by the spermatocyte specific factor, RFX2 (6). The regulatory elements for Soggy-1 lie very close to those of TEAD-2, a transcription factor that is expressed very early in development (4). Soggy-1 and TEAD-2 are co-expressed in preimplantation embryos and embryonic stem cells, but differentiated cells express only one or the other (4, 5). During development, Soggy-1 is first detectable at the onset of sexual differentiation (4). Soggy-1 is localized to the acrosome in developing spermatocytes and mature spermatozoa (3, 4). Soggy-1, as expressed in developing mouse spermatocytes, is a 34 kDa N-glycosylated protein. This glycosylation is not present on Soggy-1 in mature spermatozoa, although the apparent molecular weight suggests the presence of some post-translational modification (3). Two shorter forms of Soggy-1 have been described that result from the use of internal methionine residues for initiation (4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.