

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PDGF-C in direct ELISAs. In direct ELISAs, 100% cross-reactivity with recombinant mouse PDGF-CC is observed, and no cross-reactivity with recombinant human (rh) PDGF-AA, rhPDGF-AB, rhPDGF-BB, rhPDGF-D, or recombinant rat PDGF-AB is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 619346
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	E.coli-derived recombinant human PDGF-C Val235-Gly345 Accession # Q9NRA1
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The platelet-derived growth factor (PDGF) family consists of proteins derived from four genes (PDGF-A, -B, -C, and -D) that form four disulfide-linked homodimers (PDGF-AA, -BB, -CC, and -DD) and one heterodimer (PDGF-AB) (1). These proteins regulate diverse cellular functions by binding to and inducing the homo- or hetero-dimerization of two receptor tyrosine kinases (PDGF R α and R β). Within the PDGF family, PDGF-C and PDGF-D constitute a subgroup that shares similar structural organization (2, 3). Both proteins are secreted as inactive homodimeric latent growth factors. Each monomer has two distinct protein domains: an N-terminal CUB domain; and a C-terminal PDGF/VEGF homology domain that shares 27-35% sequence identity with the corresponding regions of other PDGF family members. An 80-90 amino acid residue hinge region connects the two domains. Sequential removal of the CUB domains in the homodimeric latent growth factor by extracellular proteolytic cleavage at the hinge region is required to release the bioactive PDGF/VEGF homology domain(1). Twelve cysteine residues are found within the PDGF/VEGF homology domain of PDGF-C, including the characteristic eight invariant cysteine residues involved in inter- and intra-chains disulfide-bonds needed for the formation of the cysteine-knot structure. Bioactive PDGF-CC binds with high-affinity to PDGF R α but not PDGF R β and activates PDGF R α homodimerization (1). PDGF-CC has also been shown to activate PDGF R α heterodimers (1). PDGF-CC is expressed in multiple embryonic and adult cell types and tissues. During embryonic development, PDGF-CC is involved in ductal morphogenesis (4). PDGF-CC is a potent angiogenic factor that stimulates vessel growth in the mouse cornea pocket assay and in the CAM assay (5). It stimulates coronary artery smooth muscle cell proliferation and may play an important role in cardiovascular development and function (6). PDGF-CC is also expressed in many tumors and tumor cell lines and has a causative role in tumorigenesis (7). Mature human and mouse PDGF-C share 93.7% amino acid sequence identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.