

## DESCRIPTION

|                                                                                                                                                                                                         |                                                                                                                                                                                                                                                      |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b>                                                                                                                                                                               | Human                                                                                                                                                                                                                                                |
| <b>Specificity</b>                                                                                                                                                                                      | Detects human Kininogen Light Chain in direct ELISAs and Western blots. In Western blots, it reacts with the full length (amino acids 19-644) and two forms of the light chain starting at amino acids 390 and 438, respectively, and does not react |
| <b>Source</b>                                                                                                                                                                                           | Monoclonal Mouse IgG <sub>2B</sub> Clone # 236006                                                                                                                                                                                                    |
| <b>Purification</b>                                                                                                                                                                                     | Protein A or G purified from hybridoma culture supernatant                                                                                                                                                                                           |
| <b>Immunogen</b>                                                                                                                                                                                        | Mouse myeloma cell line NS0-derived recombinant human Kininogen<br>Gln19-Ser644 (Ile581Thr) (predicted)<br>Accession # P01042.2                                                                                                                      |
| <b>Conjugate</b>                                                                                                                                                                                        | Alexa Fluor 488<br>Excitation Wavelength: 488 nm<br>Emission Wavelength: 515-545 nm                                                                                                                                                                  |
| <b>Formulation</b>                                                                                                                                                                                      | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide                                                                                                                                                                                         |
| *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |                                                                                                                                                                                                                                                      |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

|                            |                                                                        |
|----------------------------|------------------------------------------------------------------------|
| <b>Western Blot</b>        | Optimal dilution of this antibody should be experimentally determined. |
| <b>Immunoprecipitation</b> | Optimal dilution of this antibody should be experimentally determined. |

## PREPARATION AND STORAGE

|                                |                                                                                                                   |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

## BACKGROUND

Kininogen, also known as α<sub>2</sub>-thiol proteinase inhibitor, is a multi-function protein. There are two alternatively spliced forms, designated as the high molecular weight (HMW) and low MW (LMW) forms (1). The HMW form is synthesized as a 644 amino acid (aa) precursor with a signal peptide (aa 1-18). The mature chain (aa 19-644) is further processed into the heavy (aa 19-380) and the light (aa 390-644) chains. The active peptide bradykinin (aa 381-389) is released, which has a variety of functions including muscle contraction, hypotension and inflammation. The heavy chain consists of three cystatin-like domains, which are responsible for inhibiting cysteine proteases. The light chain consists of a His-rich domain, which is associated with the clotting activity. In comparison to the HMW form, the LMW Kininogen (427 aa) has the same sequence in its heavy chain and bradykinin, but a different sequence in its light chain (aa 402-427). The LMW form is not involved in blood clotting.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.