

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Thrombospondin-2 in ELISAs. In sandwich immunoassays, no cross-reactivity or interference with recombinant human (rh) Thrombospondin-1, -4, rhMMP-2, or rhCD36 is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 230927
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Thrombospondin-2 Gly19-Ile1172 Accession # P35442
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair) Optimal dilution of this antibody should be experimentally determined.

ELISA Detection (Matched Antibody Pair) Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Thrombospondin-2 (TSP-2) is a 150 kDa calcium-binding protein that modulates cellular interactions with extracellular matrix. Thrombospondin-1 and -2 constitute subgroup A thrombospondin family members and form disulfide-linked homotrimers, whereas Thrombospondin-3, -4, and -5/COMP constitute subgroup B and form homopentamers (1-4). The human TSP-2 cDNA encodes a 1172 amino acid (aa) precursor that includes an 18 aa signal sequence followed by an N-terminal heparin-binding domain, an oligomerization motif, one vWF-C domain, three TSP type-1 repeats, three EGF-like repeats, seven TSP type-3 repeats, and a lectin-like TSP C-terminal domain (5). Human TSP-2 shares 88-90% aa sequence identity with bovine, mouse, and rat TSP-2. Within the TSP type-3 repeats and TSP C-terminal domain, human TSP-2 shares 80% aa sequence identity with human TSP-1 and approximately 60% aa sequence identity with human TSP-3, -4, and -5/COMP. TSP-2 regulates collagen matrix formation by altering fibroblast behavior during development and in areas of tissue remodeling in the adult (6, 7). Trimerization of TSP-2 is required for the calcium-dependent cell attachment and spreading functions, while the heparin-binding domain is responsible for the destabilization of focal adhesion sites (8-10). The heparin-binding domain also mediates binding to Integrins $\alpha 3\beta 1$ and $\alpha 6\beta 1$ on microvascular endothelial cells (EC) and Integrin $\alpha 4\beta 1$ on large blood vessel EC (11, 12). A fragment of TSP-2 (heparin-binding domain, oligomerization motif, and vWF-C domain) promotes EC survival, proliferation, and chemotaxis (11). Inclusion of the three TSP type-1 domains results in a molecule that inhibits VEGF-induced EC migration and vascular tube formation (13, 14). *In vivo*, full length TSP-2 blocks tumor angiogenesis and induces vascular EC apoptosis (13, 15). HPRG functions as an apparent decoy receptor by preventing interaction of TSP-2 with CD36 on macrophages and microvasculature EC (14). TSP-2 also binds MMP-2 and facilitates MMP-2 clearance by the scavenger receptor LRP (16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.