

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SLAM/CD150 in direct ELISA and Western blots. In direct ELISAs, no cross-reactivity with recombinant human 2B4 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 123317
Purification	Protein A or G purified from ascites
Immunogen	Mouse myeloma cell line NS0-derived recombinant human SLAM/CD150 Ala21-Pro237 (predicted) Accession # Q13291
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Signaling lymphocytic activation molecule (SLAM, SLAMF1; CD150) was the first identified of a family of type I transmembrane (TM) lymphocyte activating receptors. SLAM homotypic adhesion bidirectionally stimulates T and B cells. SLAM is also expressed by hematopoietic stem cells, dendritic cells and platelets and is a T cell measles virus receptor. The 70 kDa glycoprotein contains a 216 amino acid (aa) extracellular domain (ECD) with one C2 type and one V type Ig-like domain, a 20 aa TM sequence and a 76 aa SH2-binding cytoplasmic domain. One splice variant has a shorter cytoplasmic tail and another lacks the TM sequence and is secreted. Human and mouse SLAM ECD share 60% aa identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.