

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse WISP-1/CCN4 in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody shows approximately 10-30% cross-reactivity with recombinant human (rh) WISP-1 but no cross-reactivity with rhCTGF or rmNOV.
Source	Monoclonal Rat IgG _{2B} Clone # 214203
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse WISP-1/CCN4 Leu21-Asn367 Accession # O54775
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Mouse WISP-1 (Wnt-induced secreted protein-1; also named CNN4) is a 40 kDa, secreted, heparin-binding glycoprotein that is a member of the CCN (or CTGF/Cyr61/Nov) cysteine-rich protein family (1- 6). It is synthesized as a 367 amino acid (aa) precursor that contains a series of structural homology modules. Following a 22 aa signal sequence, there is a 68 aa IGF1-like domain (aa 53-120), a 66 aa von Willebrand factor type C (VWC) module (aa 121-186), a 41 aa TSP type I domain (aa 220-260) and a 75 aa, C-terminal cysteine knot motif (aa 273-347) (1, 6). The VWC module is associated with protein-protein interaction, the TSP domain binds sulfated glycoconjugates, and the cysteine knot mediates dimerization and receptor binding (4). It is likely that WISP-1 normally circulates as an 80 kDa homodimer (2). Multiple splice forms are reported for human WISP-1, but none for rodent (7-9). Mature mouse WISP-1 is 98%, 84%, and 85% aa identical to rat, canine, and human WISP-1, respectively. WISP-1 is expressed by fibroblasts and osteoblasts, and may contribute to fracture healing by promoting bone cell formation (1, 10, 11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.