

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Pro-HGF Activator in Western blots. In Western blots, only the HGFA precursor is recognized. In Western blots, no cross-reactivity with the active form of recombinant mouse HGFA, recombinant human (rh) HGFA, or rhKallikreins 3, 5, and 11 is
Source	Monoclonal Rat IgG _{2A} Clone # 198811
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant mouse Pro-HGF Activator Gln35-Ser653 (predicted) Accession # Q9R098
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Pro-HGF-A (Hepatocyte growth factor activator protein) is a 96 kDa member of the PA-FXII-HGFA family of enzymes. It is secreted by hepatocytes and is found in plasma at approximately 40 nM concentration. Other cells selectively reported to express HGF-A include astrocytes, endothelium, fibroblasts and macrophages. Upon activation, HGF-A is known to activate both proHGF and proMSP via proteolytic cleavage. The circulating 96 kDa form of mouse HGF-A represents an inactive 619 amino acid (aa) proform (aa 35-653). Within this proform exists one FN type II domain (aa 100-147), an EGF-like domain (aa 157-195), an FN type III domain (aa 197-237), a second EGF-like domain (aa 238-276), a kringle domain (aa 283-364) and a functional peptidase S1 domain (aa 406-644). Activation of HGF-A typically occurs with a thrombin-mediated cleavage between Arg405-Ile406 and a KLK cleavage between Arg369-Val370. This creates a 34 kDa heterodimeric active product that contains a 2 kDa N-terminus (aa 370-405) disulfide linked to a 32 kDa C-terminus (aa 406-653). Once activated, the 34 kDa heterodimer binds to cell surface HAI-1, rendering it unavailable to circulating HGF. When needed, this HGF-A:HAI-1 complex is released, HGF-A dissociates from HAI-1, and HGF-A is free to activate proHGF. Over aa 35-653, mouse HGF-A shares 92% and 82% aa identity with rat and human HGF-A, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.