

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Fetuin B in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) Cystatin A, B, C, D, E/M, S, SA, SN, rhFetuin A, rhHPRG, or rhKininogen is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 212621
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Fetuin B Met19-Pro382 Accession # Q9UGM5
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fetuin is a member of the cystatin superfamily of cysteine protease inhibitors (1-3). Additional members of this superfamily are kininogen and histidine-rich glycoprotein. Fetuin A and B are two known members of the fetuin family. Hepatocytes are believed to be the principal cellular source, but other cell types also express it (4, 5). Fetuin A, also known as α₂-Heremans-Schmid glycoprotein, is an inhibitor of basic calcium phosphate precipitation and a negative acute-phase protein (6, 7). Normal circulating levels of Fetuin A in adults (300-600 µg/mL) fall significantly (30-50%) during injury and infection (7). Fetuin A and B display similarities and differences in their characteristics. Fetuin B exhibits reduction of calcification, while both mRNA levels were down-regulated during the acute phase in inflammation-induced rats (4). However, they share only 20% amino acid sequence identity (2). The amounts of Fetuin B in human serum, unlike Fetuin A, vary with gender and are higher in females than in males (4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.