

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD11c.
Source	Monoclonal Mouse IgG ₁ Clone # ICRF 3.9
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Rheumatoid synovial fluid cells and human monocyte-derived fibronectin
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human whole blood monocytes

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CD11c, also known as Integrin αX, is an approximately 150 kDa type I transmembrane glycoprotein that heterodimerizes with Integrin β2, also known as CD18. The CD11c/CD18 complex, also called CR4 (complement receptor type 4) is expressed on macrophages, dendritic cells and hairy cell leukemias, with lower amounts on other myeloid cells and activated B, NK and some cytotoxic T cells. It binds fibrinogen and has been reported to be a receptor for complement component iC3b (1-3). The human CD11c extracellular domain (amino acids 20-1107) shares 70 - 76% amino acid sequence identity with mouse, rat and canine CD11c.

References:

- Hogg, N. *et al.* (1986) *Eur. J. Immunol.* **16**(3):240.
- Knapp, W.B. *et al.* eds. (1989) *Leukocyte Typing IV: White Cell Differentiation Antigens*, Oxford University Press, New York.
- Stacker, S.A. and T.A. Springer, *J. Immunol.* (1991) **146**(2):648.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.