

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Integrin β 1/CD29 in flow cytometry.
Source	Monoclonal Mouse IgG ₁ Clone # P5D2
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human skin keratinocytes
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

The Integrin β 1 subunit, also known as CD29, associates with at least ten different Integrin α subunits. It regulates not only its multiple ligands, but activates a signaling cascade in its expressing cells. CD29 is proposed to play a role in cell adhesion, apoptosis, and differentiation (1,2). Over amino acids (aa) 21-728, human and mouse share 92% aa sequence identity.

References:

1. Barkan, D. and A.F. Chambers (2011) Clin. Cancer Res. **17**:7219.
2. Humphries, M.J. (2000) Biochem. Soc. Trans. **28**:311.