

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse SIGNR1/CD209b in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse SIGNR4 is observed.
Source	Monoclonal Rat IgG _{2B} Clone # 258829
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse SIGNR1/CD209b Ser76-Gly325 Accession # Q8CJ91
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SIGNR1 belongs to the family of C-type lectins that participate in innate immune responses by binding and clearing pathogens (1-3). SIGNR1 is one of four mouse proteins, encoded by separate genes, with significant homology to DC-SIGN and DC-SIGNR (4). Mouse SIGNR1 consists of a 53 amino acid (aa) cytoplasmic domain, a 20 aa transmembrane segment, and a 252 aa extracellular domain that contains a juxtamembrane neck region and a carbohydrate recognition domain (CRD) (4, 5). SIGNR1 is a 50kDa type II protein that forms high molecular weight aggregates as has been shown for DC-SIGN and DC-SIGNR (6, 7). The mouse SIGNR1 mRNA is extensively spliced, giving rise to multiple isoforms in addition to variability in the 5' and 3' untranslated regions. The α isoform is known as SIGNR1. The β and γ isoforms lack the transmembrane segment, with the γ isoform also showing a 21 aa insertion in the CRD. The δ isoform lacks almost the entire CRD (5). Functional differences between the isoforms have not been described. Within the CRD, mouse SIGNR1 shares 67%-70% aa sequence identity with mouse DC-SIGN, SIGNR2, 3, and 4, and 64%-68% aa sequence identity with human DC-SIGN, human L-SIGN, and rat DC-SIGN. The CRD binds mannose, GlcNAc, and fucose structures on some bacterial strains, mycobacteria, and yeast (8). SIGNR1 is expressed on macrophages found in the splenic marginal zone, lymph node medulla, and peritoneum (6, 9, 10). It plays a dominant role in the clearance of circulating *S. pneumoniae* (11-13), enhances TLR4 signaling (14), and cooperates with Dectin-1 in the nonopsonic clearance of fungal pathogens (9, 15). SIGNR1 also binds mouse ICAM-2, human ICAM-2 and -3, and HIV gp120 (10).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.