

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Integrin α 5/CD49e in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse Integrin α 5, recombinant human (rh) Integrin α 1, 2, 3, 4, 6, 7x2, 8, 9, 10, 11, M, X, D, or rhIntegrin β 1 is
Source	Monoclonal Mouse IgG _{2A} Clone # 612557
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Integrin α 5/CD49e Phe42-Tyr995 Accession # P08648
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Integrin α 5, also known as CD49e and vitronectin receptor subunit α , is a 140-150 kDa integrin alpha chain that forms dimers with at least five beta chains including β 1, 3, 5, 6, and 8. It is a 1018 amino acid (aa) residue type I membrane protein with a large (962 aa) extracellular domain (ECD) and a short (32 aa) cytoplasmic tail. The N-terminal region of Integrin α 5, which is important for ligand binding, contains seven FG-GAP (phenylalanyl-glycyl and glycyl-alanyl-prolyl) consensus repeats that fold into a β -propeller domain. Furin cleavage of the Integrin α 5 ECD occurs after Gly 889, generating a disulfide-linked, heteromeric subunit Integrin α 5 chain. Integrin α 5-containing integrins bind multiple ECM molecules, including Vitronectin, Osteopontin, MMP-2, and TSP. The ECD of human Integrin α 5 shares 92% aa sequence identity with mouse Integrin α 5 ECD.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.