

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse FGF R5/FGFRL1 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) FGF R1α (IIIb), 1α (IIIc), 1β (IIIb), 1β (IIIc), recombinant mouse (rm) FGF R2α (IIIb), 2α (IIIc), 2
Source	Monoclonal Rat IgG _{2B} Clone # 223003
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse FGF R5/FGFRL1 Ala21-Pro374 Accession # Q91V87
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fibroblast growth factor receptor 5 (FGF R5), also known as FGRL1, is a 65 kDa transmembrane member of the FGF receptor family (1). Mature mouse FGF R5 consists of a 354 amino acid (aa) extracellular domain (ECD) with three immunoglobulin-like domains, a 21 aa transmembrane segment, and a 134 aa cytoplasmic domain (2). FGF R5 is distinct from other FGF receptor family members in that it does not contain a cytoplasmic tyrosine kinase domain (2, 3). Within the ECD, mouse FGF R5 shares 94% and 98% aa sequence identity with human and rat FGF R5, respectively. Alternate splicing results in an isoform that lacks the first Ig-like domain (2). FGF R5 is widely expressed, and expression of two species of mRNA in cartilage and pancreas may indicate the presence of both splice forms (2-6). Both the full length and truncated isoforms of FGF R5 bind FGF basic; the full length form additionally binds to heparin (2, 6). FGF R5 may function as a decoy receptor by binding FGF but not transducing its mitogenic signals (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.